



Contents lists available at ScienceDirect

Journal of Mathematical Analysis and Applications

www.elsevier.com/locate/jmaa



An overdetermined problem for harmonic functions

Robert Dalmasso

Laboratoire Jean Kuntzmann, Equipe EDP, 51 rue des Mathématiques, Domaine Universitaire, BP 53, 38041 Grenoble Cedex 9, France

ARTICLE INFO

Article history:

Received 9 May 2011

Available online 22 July 2011

Submitted by V. Radulescu

Keywords:

Overdetermined problem

Harmonic functions

ABSTRACT

Let $\Omega \subset \mathbb{R}^n$ be a bounded connected open set with connected real analytic boundary. We show that, if there exist n harmonic functions satisfying some appropriate boundary conditions, then Ω is a ball.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

The purpose of this paper is to prove the following theorem.

Theorem 1. Let $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) be a bounded connected open set with connected real analytic boundary $\partial\Omega$. We denote by $\nu = (\nu_1, \dots, \nu_n)$ the outward unit normal to $\partial\Omega$ and by H the mean curvature of $\partial\Omega$. Suppose that there exist a constant $c \in \mathbb{R} \setminus \{0\}$ and n harmonic functions in Ω h_1, \dots, h_n satisfying

$$h_j = c\nu_j \quad \text{on } \partial\Omega, \quad (1)$$

and

$$\frac{\partial h_j}{\partial \nu} = -(1 + c(n-1)H)\nu_j \quad \text{on } \partial\Omega, \quad (2)$$

for $j = 1, \dots, n$. Then Ω is a ball.

Notice that assumptions (1) and (2) imply that both $h = (h_1, \dots, h_n)$ and $\partial h / \partial \nu = (\partial h_1 / \partial \nu, \dots, \partial h_n / \partial \nu)$ are parallel to ν .

Remark 1. Clearly, for each $j \in \{1, \dots, n\}$, h_j extends to an analytic function on a neighborhood of $\partial\Omega$.

As in [1] we have the following corollary which gives a characterization of open balls in \mathbb{R}^n by means of integral identities involving harmonic functions.

Corollary. Let $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) be a bounded connected open set with connected real analytic boundary $\partial\Omega$. We denote by $\nu = (\nu_1, \dots, \nu_n)$ the outward unit normal to $\partial\Omega$ and by H the mean curvature of $\partial\Omega$. If

E-mail address: robert.dalmasso@imag.fr.

$$C \int_{\partial\Omega} \frac{\partial h}{\partial \nu} v_j ds = - \int_{\partial\Omega} (1 + C(n-1)H) h v_j ds, \quad \forall j \in \{1, \dots, n\}, \quad (3)$$

for some constant $C \in \mathbb{R}$ and for every $h \in C^2(\Omega) \cap C^1(\overline{\Omega})$ such that $\Delta h = 0$ in Ω , then Ω is a ball.

Remark 2. Notice that necessarily $C \neq 0$ in (3). Indeed let h be such that $\Delta h = 0$ in Ω and $h = v_j$ on $\partial\Omega$ for some $j \in \{1, \dots, n\}$. Then (3) implies that $C \neq 0$.

Remark 3. Suppose that there exist a constant $c \in \mathbb{R} \setminus \{0\}$ and n harmonic functions h_1, \dots, h_n satisfying (1)–(2). Using Green's identity we get

$$\int_{\partial\Omega} h_j \frac{\partial h}{\partial \nu} ds = \int_{\partial\Omega} h \frac{\partial h_j}{\partial \nu} ds, \quad j = 1, \dots, n,$$

for every harmonic function $h \in C^2(\Omega) \cap C^1(\overline{\Omega})$. Then (3) holds with $C = c$.

We shall need the following result [8].

Theorem 2. Let $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) be a bounded connected open set with C^2 boundary. Suppose that there exists $u \in C^2(\overline{\Omega})$ such that

$$\begin{aligned} \Delta u + 1 &= 0 \quad \text{in } \Omega, \\ u &= 0, \quad \frac{\partial u}{\partial \nu} = d \quad \text{on } \partial\Omega, \end{aligned}$$

where d denotes some constant. Then Ω is a ball of center x_0 and radius $R = -dn$ and $u(x) = (R^2 - |x - x_0|^2)/2n$ for $x \in \Omega$.

Remark 4. Let u be as in Theorem 2. For each $j \in \{1, \dots, n\}$, $\partial u / \partial x_j$ is harmonic in Ω . Since $u = 0$ on $\partial\Omega$, we have $\partial u / \partial x_j = (\partial u / \partial \nu) v_j = d v_j$ on $\partial\Omega$ for $j = 1, \dots, n$. Now, arguing as in the proof of Lemma 2 below, we easily obtain

$$\frac{\partial}{\partial \nu} \left(\frac{\partial u}{\partial x_j} \right) = -(1 + d(n-1)H) v_j, \quad j = 1, \dots, n.$$

To the best of our knowledge Theorem 1 is the first result in this direction. It would be interesting to give a proof of Theorem 1 that does not make use of Serrin's result. Finally let us also mention that there is a huge amount of literature concerning overdetermined problems in both bounded and unbounded domains: See [2–7,9–11] and the references therein.

2. Proof of Theorem 1

$\partial\Omega$ is locally the graph of an analytic function of $n-1$ variables φ , that is, $\partial\Omega$ is locally defined by $x_n = \varphi(x_1, \dots, x_{n-1})$ and $x = (x_1, \dots, x_n) = (x', x_n) \in \Omega$ if and only if $x_n < \varphi(x_1, \dots, x_{n-1}) = \varphi(x')$. Then the outward unit normal ν and the mean curvature H at $x = (x', \varphi(x')) \in \partial\Omega$ are given by

$$\nu(x) = \frac{1}{\sqrt{1 + |\nabla \varphi(x')|^2}} (-\nabla \varphi(x'), 1),$$

and

$$H(x) = \frac{1}{n-1} \sum_{j=1}^{n-1} \frac{\partial}{\partial x_j} (v_j(x)).$$

We shall use repeatedly the following lemma.

Lemma 1. We have

$$\sum_{j=1}^{n-1} v_j \frac{\partial}{\partial x_j} (v_n) = v_n ((n-1)H + v_n \Delta \varphi),$$

and

$$\sum_{j=1}^{n-1} v_j \frac{\partial}{\partial x_j} (v_p) = -\frac{1}{v_n} \frac{\partial}{\partial x_p} (v_n) + v_p ((n-1)H + v_n \Delta \varphi)$$

for $1 \leq p \leq n-1$.

Proof. Let $p \in \{1, \dots, n-1\}$. We have

$$\begin{aligned} \sum_{j=1}^{n-1} v_j \frac{\partial}{\partial x_j} (v_p) &= \sum_{j=1}^{n-1} v_j \left(-v_n \frac{\partial^2 \varphi}{\partial x_j \partial x_p} + v_n^3 \frac{\partial \varphi}{\partial x_p} \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial \varphi}{\partial x_k} \right) \\ &= v_n^2 \sum_{j=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_p} \frac{\partial \varphi}{\partial x_j} + v_p v_n^3 \sum_{k=1}^{n-1} \frac{\partial \varphi}{\partial x_k} \left(\sum_{j=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial \varphi}{\partial x_j} \right) \\ &= -\frac{1}{v_n} \frac{\partial}{\partial x_p} (v_n) + v_p \sum_{k=1}^{n-1} \left(\frac{\partial}{\partial x_k} (v_k) + v_n \frac{\partial^2 \varphi}{\partial x_k^2} \right) \\ &= -\frac{1}{v_n} \frac{\partial}{\partial x_p} (v_n) + v_p ((n-1)H + v_n \Delta \varphi). \end{aligned}$$

The proof is analogous when $p = n$. \square

In view of Remark 4, our first aim is to show that there exists an analytic function w satisfying

$$\frac{\partial w}{\partial x_j} = h_j \quad \text{in } \Omega, \quad j = 1, \dots, n.$$

Therefore it is natural to expect that the functions h_j satisfy Lemma 5 below. Unfortunately long and tedious calculations are required.

Lemma 2. We have

$$\frac{\partial h_j}{\partial x_k} = -(1 + 2c(n-1)H + cv_n \Delta \varphi) v_j v_k + c \frac{v_k}{v_n} \frac{\partial}{\partial x_j} (v_n) + c \frac{\partial}{\partial x_k} (v_j)$$

on $\partial \Omega$ for $1 \leq j, k \leq n$. Moreover

$$\frac{\partial h_j}{\partial x_k} = \frac{\partial h_k}{\partial x_j} \quad \text{for } 1 \leq j, k \leq n, \quad \text{and} \quad \sum_{j=1}^n \frac{\partial h_j}{\partial x_j} = -1$$

on $\partial \Omega$.

Proof. Differentiating (1) with respect to x_k for $k \in \{1, \dots, n-1\}$ and multiplying by v_n we get

$$v_n \frac{\partial h_j}{\partial x_k} - v_k \frac{\partial h_j}{\partial x_n} = cv_n \frac{\partial}{\partial x_k} (v_j), \tag{4}$$

for $1 \leq j \leq n$. Multiplying (4) by v_k and adding we obtain

$$v_n \frac{\partial h_j}{\partial v} - \frac{\partial h_j}{\partial x_n} = cv_n \sum_{k=1}^{n-1} v_k \frac{\partial}{\partial x_k} (v_j).$$

Then using (2) we deduce that

$$\frac{\partial h_j}{\partial x_n} = -(1 + c(n-1)H) v_j v_n - cv_n \sum_{k=1}^{n-1} v_k \frac{\partial}{\partial x_k} (v_j), \tag{5}$$

for $1 \leq j \leq n$. Now (4) and (5) imply that

$$\frac{\partial h_j}{\partial x_k} = -(1 + c(n-1)H) v_j v_k - cv_k \sum_{r=1}^{n-1} v_r \frac{\partial}{\partial x_r} (v_j) + c \frac{\partial}{\partial x_k} (v_j), \tag{6}$$

for $1 \leq j \leq n$ and $1 \leq k \leq n-1$. (5), (6) and Lemma 1 give the first part of the lemma. Now with the help of Lemma 1 we get

$$\begin{aligned}
\sum_{j=1}^n \frac{\partial h_j}{\partial x_j} &= -(1 + 2c(n-1)H + cv_n \Delta \varphi) + \frac{c}{v_n} \sum_{j=1}^{n-1} v_j \frac{\partial}{\partial x_j} (v_n) + c \sum_{j=1}^{n-1} \frac{\partial}{\partial x_j} (v_j) \\
&= -(1 + 2c(n-1)H + cv_n \Delta \varphi) + c((n-1)H + v_n \Delta \varphi) + c(n-1)H \\
&= -1
\end{aligned}$$

on $\partial\Omega$. Finally for $1 \leq j, k \leq n-1$ we have

$$\begin{aligned}
\frac{v_k}{v_n} \frac{\partial}{\partial x_j} (v_n) + \frac{\partial}{\partial x_k} (v_j) &= -v_k v_n^2 \sum_{i=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_j} \frac{\partial \varphi}{\partial x_i} - v_n \frac{\partial^2 \varphi}{\partial x_j \partial x_k} + v_n^3 \frac{\partial \varphi}{\partial x_j} \sum_{i=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_i} \\
&= -v_k v_n^2 \sum_{i=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_j} \frac{\partial \varphi}{\partial x_i} - v_n \frac{\partial^2 \varphi}{\partial x_j \partial x_k} - v_j v_n^2 \sum_{i=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_i} \\
&= \frac{v_j}{v_n} \frac{\partial}{\partial x_k} (v_n) + \frac{\partial}{\partial x_j} (v_k),
\end{aligned}$$

and we conclude with the first part of the lemma. \square

Lemma 3. Define $a_{jk} : \partial\Omega \rightarrow \mathbb{R}$ by $a_{jk} = \partial h_j / \partial x_k$. We have

$$\frac{v_k}{\partial x_k \partial x_p} \frac{\partial^2 h_j}{\partial x_k \partial x_p} = -v_k v_p \left(v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) + \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) \right) + v_n \frac{\partial}{\partial x_p} (a_{jk}) + v_p \frac{\partial}{\partial x_k} (a_{jn})$$

on $\partial\Omega$ for $1 \leq j, k, p \leq n$.

Proof. Differentiating a_{jk} with respect to x_p for $p \in \{1, \dots, n-1\}$ and multiplying by v_n we get

$$v_n \frac{\partial^2 h_j}{\partial x_k \partial x_p} - v_p \frac{\partial^2 h_j}{\partial x_k \partial x_n} = v_n \frac{\partial}{\partial x_p} (a_{jk}), \quad (7)$$

for $1 \leq j, k \leq n$. Let $k = p$ in (7). Adding and using Lemma 2 we obtain

$$\frac{\partial}{\partial v} \left(\frac{\partial h_j}{\partial x_n} \right) = v_n \Delta h_j - v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) = -v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}), \quad (8)$$

for $1 \leq j \leq n$. Multiplying (7) by v_p and adding we obtain

$$\frac{\partial^2 h_j}{\partial x_k \partial x_n} = v_n \frac{\partial}{\partial v} \left(\frac{\partial h_j}{\partial x_k} \right) - v_n \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jk}), \quad (9)$$

for $1 \leq j, k \leq n$. From (8) and (9) we deduce that

$$\frac{\partial^2 h_j}{\partial x_n^2} = -v_n \left(v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) + \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) \right) \quad (10)$$

for $1 \leq j \leq n$. From (7) and (10) we get

$$\begin{aligned}
v_n \frac{\partial^2 h_j}{\partial x_p \partial x_n} &= v_p \frac{\partial^2 h_j}{\partial x_n^2} + v_n \frac{\partial}{\partial x_p} (a_{jn}) \\
&= -v_p v_n \left(v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) + \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) \right) + v_n \frac{\partial}{\partial x_p} (a_{jn}),
\end{aligned}$$

for $1 \leq j \leq n$ and $1 \leq p \leq n-1$, hence

$$\frac{\partial^2 h_j}{\partial x_p \partial x_n} = -v_p \left(v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) + \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) \right) + \frac{\partial}{\partial x_p} (a_{jn}), \quad (11)$$

for $1 \leq j \leq n$ and $1 \leq p \leq n-1$. Finally using (7) and (11) we have

$$\begin{aligned} v_n \frac{\partial^2 h_j}{\partial x_k \partial x_p} &= v_p \frac{\partial^2 h_j}{\partial x_k \partial x_n} + v_n \frac{\partial}{\partial x_p} (a_{jk}) \\ &= -v_k v_p \left(v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) + \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) \right) + v_p \frac{\partial}{\partial x_k} (a_{jn}) + v_n \frac{\partial}{\partial x_p} (a_{jk}), \end{aligned}$$

for $1 \leq j \leq n$ and $1 \leq k, p \leq n-1$ and the lemma is proved. \square

Lemma 4. We have

$$\frac{\partial^2 h_j}{\partial x_k \partial x_p} = \frac{\partial^2 h_k}{\partial x_j \partial x_p}$$

on $\partial\Omega$ for $1 \leq j, k, p \leq n$.

Proof. It is enough to show that

$$\frac{\partial^2 h_j}{\partial x_n^2} = \frac{\partial^2 h_n}{\partial x_j \partial x_n} \quad (12)$$

on $\partial\Omega$ for $1 \leq j \leq n-1$. Indeed assume that (12) holds. (7) and Lemma 2 imply that

$$\begin{aligned} v_n \frac{\partial^2 h_j}{\partial x_p \partial x_n} &= v_p \frac{\partial^2 h_j}{\partial x_n^2} + v_n \frac{\partial}{\partial x_p} (a_{jn}) \\ &= v_p \frac{\partial^2 h_n}{\partial x_j \partial x_n} + v_n \frac{\partial}{\partial x_p} (a_{jn}) \\ &= v_n \frac{\partial^2 h_n}{\partial x_p \partial x_j}, \end{aligned}$$

for $1 \leq j \leq n$ and $1 \leq p \leq n-1$. Therefore the lemma holds for $k = n$ and $1 \leq j, p \leq n$. Then we have

$$\frac{\partial^2 h_j}{\partial x_k \partial x_n} = \frac{\partial^2 h_n}{\partial x_j \partial x_k} = \frac{\partial^2 h_k}{\partial x_j \partial x_n}.$$

Using (7) again we get

$$\begin{aligned} v_n \frac{\partial^2 h_j}{\partial x_k \partial x_p} &= v_p \frac{\partial^2 h_j}{\partial x_k \partial x_n} + v_n \frac{\partial}{\partial x_p} (a_{jk}) = v_p \frac{\partial^2 h_n}{\partial x_j \partial x_k} + v_n \frac{\partial}{\partial x_p} (a_{jk}) \\ &= v_p \frac{\partial^2 h_k}{\partial x_j \partial x_n} + v_n \frac{\partial}{\partial x_p} (a_{jk}) = v_n \frac{\partial^2 h_k}{\partial x_j \partial x_p}, \end{aligned}$$

and the lemma is proved. Now we prove (12). In view of Lemma 3 we have to show that

$$v_n \left(v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) + \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) \right) = v_j \left(v_n \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{in}) + \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{nn}) \right) - \frac{\partial}{\partial x_j} (a_{nn}), \quad (13)$$

on $\partial\Omega$ for $1 \leq j \leq n-1$. Define

$$a = 1 + 2c(n-1)H + cv_n \Delta \varphi.$$

By Lemma 2 we have

$$a_{ij} = a_{ji} = -av_i v_j + c \frac{v_i}{v_n} \frac{\partial}{\partial x_j} (v_n) + c \frac{\partial}{\partial x_i} (v_j).$$

Then, using Lemma 1, we get

$$\begin{aligned} \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) &= -v_j \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a) - a \sum_{i=1}^{n-1} \left(v_j \frac{\partial}{\partial x_i} (v_i) + v_i \frac{\partial}{\partial x_i} (v_j) \right) \\ &\quad - c \Delta \varphi \frac{\partial}{\partial x_j} (v_n) + \frac{c}{v_n} \sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n) + c \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_j) \end{aligned}$$

$$\begin{aligned}
&= -v_j \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a) - a \left(2(n-1)H v_j - \frac{1}{v_n} \frac{\partial}{\partial x_j} (v_n) + v_n v_j \Delta \varphi \right) \\
&\quad - c \Delta \varphi \frac{\partial}{\partial x_j} (v_n) + \frac{c}{v_n} \sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n) + c \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_j),
\end{aligned}$$

and

$$\begin{aligned}
\sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) &= -v_j v_n \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a) - a \sum_{i=1}^{n-1} v_i \left(v_j \frac{\partial}{\partial x_i} (v_n) + v_n \frac{\partial}{\partial x_i} (v_j) \right) + c \sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n) \\
&= -v_j v_n \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a) + c \sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n) - a \left(2v_j v_n ((n-1)H + v_n \Delta \varphi) - \frac{\partial}{\partial x_j} (v_n) \right),
\end{aligned}$$

from which we deduce that

$$\begin{aligned}
v_n v_j \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{in}) - v_n^2 \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (a_{ij}) &= -a v_n \frac{\partial}{\partial x_j} (v_n) + c v_n^2 \Delta \varphi \frac{\partial}{\partial x_j} (v_n) - c v_n \sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n) \\
&\quad - c v_n^2 \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_j) + c v_n v_j \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_n),
\end{aligned}$$

and

$$v_j \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{nn}) - v_n \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (a_{jn}) = -a v_n \frac{\partial}{\partial x_j} (v_n) - c v_n \sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n).$$

Define

$$A = c v_n^2 \Delta \varphi \frac{\partial}{\partial x_j} (v_n) + v_n^2 \frac{\partial}{\partial x_j} (a) - 2c v_n \sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n) - c v_n^2 \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_j) + c v_n v_j \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_n).$$

Since

$$\frac{\partial}{\partial x_j} (a_{nn}) = -a \frac{\partial}{\partial x_j} (v_n^2) - v_n^2 \frac{\partial}{\partial x_j} (a),$$

we deduce that (13) is equivalent to $A = 0$. Let $1 \leq i, j \leq n-1$. We have

$$\frac{\partial}{\partial x_i} (v_n) = -v_n^3 \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_k},$$

and

$$\frac{\partial}{\partial x_i} (v_j) = -v_n \frac{\partial^2 \varphi}{\partial x_i \partial x_j} + v_n^3 \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_k} \frac{\partial \varphi}{\partial x_j}.$$

Then

$$\begin{aligned}
\frac{\partial^2}{\partial x_i^2} (v_n) &= \frac{3}{v_n} \left(\frac{\partial}{\partial x_i} (v_n) \right)^2 - v_n^3 \sum_{k=1}^{n-1} \left(\frac{\partial^3 \varphi}{\partial x_i^2 \partial x_k} \frac{\partial \varphi}{\partial x_k} + \left(\frac{\partial^2 \varphi}{\partial x_i \partial x_k} \right)^2 \right), \\
\frac{\partial^2}{\partial x_i \partial x_j} (v_n) &= \frac{3}{v_n} \frac{\partial}{\partial x_i} (v_n) \frac{\partial}{\partial x_j} (v_n) - v_n^3 \sum_{k=1}^{n-1} \left(\frac{\partial^3 \varphi}{\partial x_i \partial x_j \partial x_k} \frac{\partial \varphi}{\partial x_k} + \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \right)
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial^2}{\partial x_i^2} (v_j) &= -\frac{\partial^2 \varphi}{\partial x_i \partial x_j} \frac{\partial}{\partial x_i} (v_n) - v_n \frac{\partial^3 \varphi}{\partial x_i^2 \partial x_j} + \frac{3}{v_n^2} v_j \left(\frac{\partial}{\partial x_i} (v_n) \right)^2 \\
&\quad + v_n^3 \sum_{k=1}^{n-1} \left(\left(\frac{\partial^3 \varphi}{\partial x_i^2 \partial x_k} \frac{\partial \varphi}{\partial x_k} + \left(\frac{\partial^2 \varphi}{\partial x_i \partial x_k} \right)^2 \right) \frac{\partial \varphi}{\partial x_j} + \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \right).
\end{aligned}$$

We deduce that

$$\begin{aligned}\sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_n) &= \frac{3}{v_n} \sum_{i=1}^{n-1} \left(\frac{\partial}{\partial x_i} (v_n) \right)^2 + v_n^2 \sum_{k=1}^{n-1} v_k \frac{\partial \Delta \varphi}{\partial x_k} - v_n^3 \sum_{i,k=1}^{n-1} \left(\frac{\partial^2 \varphi}{\partial x_i \partial x_k} \right)^2, \\ \sum_{i=1}^{n-1} \frac{\partial^2}{\partial x_i^2} (v_j) &= -2 \sum_{i=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_j} \frac{\partial}{\partial x_i} (v_n) - v_n \frac{\partial \Delta \varphi}{\partial x_j} + \frac{3}{v_n^2} v_j \sum_{i=1}^{n-1} \left(\frac{\partial}{\partial x_i} (v_n) \right)^2 \\ &\quad + v_n v_j \sum_{k=1}^{n-1} v_k \frac{\partial \Delta \varphi}{\partial x_k} - v_n^2 v_j \sum_{i,k=1}^{n-1} \left(\frac{\partial^2 \varphi}{\partial x_i \partial x_k} \right)^2,\end{aligned}$$

and with the help of Lemma 1

$$\sum_{i=1}^{n-1} v_i \frac{\partial^2}{\partial x_i \partial x_j} (v_n) = 3((n-1)H + v_n \Delta \varphi) \frac{\partial}{\partial x_j} (v_n) + v_n^2 \sum_{i,k=1}^{n-1} \frac{\partial^3 \varphi}{\partial x_i \partial x_j \partial x_k} v_i v_k - v_n \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial}{\partial x_k} (v_n).$$

Therefore

$$\begin{aligned}A &= -4c v_n^2 \Delta \varphi \frac{\partial}{\partial x_j} (v_n) - 6c v_n (n-1) H \frac{\partial}{\partial x_j} (v_n) - 2c v_n^3 \sum_{i,k=1}^{n-1} \frac{\partial^3 \varphi}{\partial x_i \partial x_j \partial x_k} v_i v_k \\ &\quad + 4c v_n^2 \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial}{\partial x_k} (v_n) + 2c v_n^3 \frac{\partial \Delta \varphi}{\partial x_j} + 2c(n-1) v_n^2 \frac{\partial}{\partial x_j} (H).\end{aligned}$$

Now we have

$$\begin{aligned}(n-1)H &= \sum_{i=1}^{n-1} \frac{\partial}{\partial x_i} (v_i) = \sum_{i=1}^{n-1} \left(-\frac{\partial^2 \varphi}{\partial x_i^2} v_n + v_n^3 \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_k} \frac{\partial \varphi}{\partial x_i} \right) \\ &= -v_n \Delta \varphi + v_n^3 \sum_{i,k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_k} \frac{\partial \varphi}{\partial x_i},\end{aligned}$$

hence

$$\begin{aligned}(n-1) \frac{\partial}{\partial x_j} (H) &= -\Delta \varphi \frac{\partial}{\partial x_j} (v_n) - v_n \frac{\partial \Delta \varphi}{\partial x_j} + 3v_n^2 \frac{\partial}{\partial x_j} (v_n) \sum_{i,k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_k} \frac{\partial \varphi}{\partial x_i} \\ &\quad + v_n^3 \sum_{i,k=1}^{n-1} \left(\frac{\partial^3 \varphi}{\partial x_i \partial x_j \partial x_k} \frac{\partial \varphi}{\partial x_k} \frac{\partial \varphi}{\partial x_i} + \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial \varphi}{\partial x_i} + \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial^2 \varphi}{\partial x_i \partial x_j} \frac{\partial \varphi}{\partial x_k} \right) \\ &= -\Delta \varphi \frac{\partial}{\partial x_j} (v_n) - v_n \frac{\partial \Delta \varphi}{\partial x_j} - 3v_n \frac{\partial}{\partial x_j} (v_n) \sum_{i,k=1}^{n-1} v_i \frac{\partial^2 \varphi}{\partial x_i \partial x_k} \frac{\partial \varphi}{\partial x_k} \\ &\quad + v_n \sum_{i,k=1}^{n-1} \frac{\partial^3 \varphi}{\partial x_i \partial x_j \partial x_k} v_i v_k - 2 \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial}{\partial x_k} (v_n) \\ &= -\Delta \varphi \frac{\partial}{\partial x_j} (v_n) - v_n \frac{\partial \Delta \varphi}{\partial x_j} + \frac{3}{v_n^2} \frac{\partial}{\partial x_j} (v_n) \sum_{i=1}^{n-1} v_i \frac{\partial}{\partial x_i} (v_n) \\ &\quad + v_n \sum_{i,k=1}^{n-1} \frac{\partial^3 \varphi}{\partial x_i \partial x_j \partial x_k} v_i v_k - 2 \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial}{\partial x_k} (v_n) \\ &= 2\Delta \varphi \frac{\partial}{\partial x_j} (v_n) - v_n \frac{\partial \Delta \varphi}{\partial x_j} + \frac{3}{v_n} (n-1) H \frac{\partial}{\partial x_j} (v_n) + v_n \sum_{i,k=1}^{n-1} \frac{\partial^3 \varphi}{\partial x_i \partial x_j \partial x_k} v_i v_k - 2 \sum_{k=1}^{n-1} \frac{\partial^2 \varphi}{\partial x_j \partial x_k} \frac{\partial}{\partial x_k} (v_n)\end{aligned}$$

where we have again used Lemma 1. Now we easily verify that $A = 0$ and (13) is proved. The proof of the lemma is complete. \square

Lemma 5. Let $\alpha \in \mathbb{N}^n$. We have

$$\frac{\partial^{|\alpha|+1} h_j}{\partial x_k \partial x^\alpha} = \frac{\partial^{|\alpha|+1} h_k}{\partial x_j \partial x^\alpha} \quad (14)$$

on $\partial\Omega$ for $1 \leq j, k \leq n$.

Proof. Lemmas 3 and 4 imply that (14) holds for $|\alpha| \leq 1$. Now suppose that (14) holds for $|\alpha| = m \geq 1$. Then for all $\beta \in \mathbb{N}^n$ such that $|\beta| = m - 1$ and for all $p \in \{1, \dots, n\}$ we have

$$\frac{\partial^{|\beta|+2} h_j}{\partial x_k \partial x_p \partial x^\beta} = \frac{\partial^{|\beta|+2} h_k}{\partial x_j \partial x_p \partial x^\beta} \quad (15)$$

on $\partial\Omega$ for $1 \leq j, k \leq n$. Differentiating (15) with respect to x_i for $i \in \{1, \dots, n-1\}$ and multiplying by v_n we get

$$v_n \frac{\partial^{|\beta|+3} h_j}{\partial x_i \partial x_k \partial x_p \partial x^\beta} - v_i \frac{\partial^{|\beta|+3} h_j}{\partial x_k \partial x_p \partial x_n \partial x^\beta} = v_n \frac{\partial^{|\beta|+3} h_k}{\partial x_i \partial x_j \partial x_p \partial x^\beta} - v_i \frac{\partial^{|\beta|+3} h_k}{\partial x_j \partial x_p \partial x_n \partial x^\beta}, \quad (16)$$

for $1 \leq j, k, p \leq n$. Let $i = p$ in (16). Adding we get

$$v_n \Delta \frac{\partial^{|\beta|+1} h_j}{\partial x_k \partial x^\beta} - \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_j}{\partial x_k \partial x_n \partial x^\beta} \right) = v_n \Delta \frac{\partial^{|\beta|+1} h_k}{\partial x_j \partial x^\beta} - \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_k}{\partial x_j \partial x_n \partial x^\beta} \right).$$

Therefore we have

$$\frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_j}{\partial x_k \partial x_n \partial x^\beta} \right) = \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_k}{\partial x_j \partial x_n \partial x^\beta} \right), \quad (17)$$

for $1 \leq j, k \leq n$. Multiplying (16) by v_i and adding we obtain

$$v_n \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_j}{\partial x_k \partial x_p \partial x^\beta} \right) - \frac{\partial^{|\beta|+3} h_j}{\partial x_k \partial x_p \partial x_n \partial x^\beta} = v_n \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_k}{\partial x_j \partial x_p \partial x^\beta} \right) - \frac{\partial^{|\beta|+3} h_k}{\partial x_j \partial x_p \partial x_n \partial x^\beta}, \quad (18)$$

for $1 \leq j, k, p \leq n$. Multiplying (16) by v_p and adding we obtain

$$v_n \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_j}{\partial x_i \partial x_k \partial x^\beta} \right) - v_i \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_j}{\partial x_k \partial x_n \partial x^\beta} \right) = v_n \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_k}{\partial x_i \partial x_j \partial x^\beta} \right) - v_i \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_k}{\partial x_j \partial x_n \partial x^\beta} \right), \quad (19)$$

for $1 \leq j, k \leq n$ and $1 \leq i \leq n-1$. (17) and (19) imply that

$$\frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_j}{\partial x_i \partial x_k \partial x^\beta} \right) = \frac{\partial}{\partial v} \left(\frac{\partial^{|\beta|+2} h_k}{\partial x_i \partial x_j \partial x^\beta} \right), \quad (20)$$

for $1 \leq i \leq n-1$ and $1 \leq j, k \leq n$. Then (17), (18) and (20) give

$$\frac{\partial^{|\beta|+3} h_j}{\partial x_k \partial x_p \partial x_n \partial x^\beta} = \frac{\partial^{|\beta|+3} h_k}{\partial x_j \partial x_p \partial x_n \partial x^\beta}, \quad (21)$$

for $1 \leq j, k, p \leq n$. Now (18) and (20) imply

$$\frac{\partial^{|\beta|+3} h_j}{\partial x_i \partial x_k \partial x_p \partial x^\beta} = \frac{\partial^{|\beta|+3} h_k}{\partial x_i \partial x_j \partial x_p \partial x^\beta}, \quad (22)$$

for $1 \leq j, k, p \leq n$ and $1 \leq i \leq n-1$. The lemma follows from (21) and (22). \square

Lemma 6. Let $y \in \partial\Omega$. There exist $r > 0$ and a function v analytic in the open ball $B(y, r)$ of \mathbb{R}^n such that

$$\frac{\partial v}{\partial x_j}(x) = h_j(x), \quad \forall x \in B(y, r), \quad j = 1, \dots, n.$$

Proof. By Remark 1 there exists an open ball $B(y, r)$ in \mathbb{R}^n such that

$$h_j(x) = \sum_{\alpha \geq 0} \frac{1}{\alpha!} \frac{\partial^{|\alpha|} h_j}{\partial x^\alpha}(y) (x - y)^\alpha$$

for $x \in B(y, r)$ and $j = 1, \dots, n$. Let $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$. We define $v_\alpha \in \mathbb{R}$ as follows: $v_0 \in \mathbb{R}$

$$v_{(\alpha_1, 0, \dots, 0)} = \frac{\partial^{\alpha_1-1} h_1}{\partial x_1^{\alpha_1-1}}(y) \quad \text{for } \alpha_1 \geq 1,$$

$$\vdots$$

$$v_{(0, \dots, 0, \alpha_n)} = \frac{\partial^{\alpha_n-1} h_n}{\partial x_n^{\alpha_n-1}}(y) \quad \text{for } \alpha_n \geq 1,$$

and, when there exist $\alpha_j, \alpha_k \geq 1$ for j and k such that $1 \leq j < k \leq n$

$$v_\alpha = \frac{\partial^{|\beta|+1} h_j}{\partial x_k \partial x^\beta}(y) = \frac{\partial^{|\beta|+1} h_k}{\partial x_j \partial x^\beta}(y),$$

where $\beta = (\alpha_1, \dots, \alpha_{j-1}, \alpha_j - 1, \alpha_{j+1}, \dots, \alpha_{k-1}, \alpha_k - 1, \alpha_{k+1}, \dots, \alpha_n)$ with the obvious modifications when $j = 1$ or $k = n$ or $k = j + 1$ and where the last equality is given by Lemma 5. Let $\varepsilon_1 = (1, 0, \dots, 0), \dots, \varepsilon_n = (0, \dots, 0, 1)$. Since

$$\begin{aligned} h_j(x) &= \sum_{\alpha \geq 0} \frac{1}{\alpha!} v_{\alpha + \varepsilon_j} (x - y)^\alpha \\ &= \frac{\partial}{\partial x_j} \left(\sum_{\alpha \geq 0} \frac{1}{\alpha!} v_\alpha (x - y)^\alpha \right), \end{aligned}$$

for $x \in B(y, r)$ and $1 \leq j \leq n$, we define an analytic function v in $B(y, r)$ by

$$v(x) = \sum_{\alpha \geq 0} \frac{1}{\alpha!} v_\alpha (x - y)^\alpha, \quad x \in B(y, r),$$

such that

$$\frac{\partial v}{\partial x_j}(x) = h_j(x), \quad x \in B(y, r), \quad j = 1, \dots, n. \quad \square$$

Lemma 7. *There exist an open connected neighborhood U of $\partial\Omega$ and a function v analytic in U such that*

$$\frac{\partial v}{\partial x_j}(x) = h_j(x), \quad \forall x \in U, \quad j = 1, \dots, n.$$

Proof. Since $\partial\Omega$ is compact, using Lemma 6, we can find $y_1, \dots, y_m \in \partial\Omega$, $r_1, \dots, r_m > 0$ and v_1, \dots, v_m analytic in $B(y_1, r_1), \dots, B(y_m, r_m)$ respectively such that

$$\partial\Omega \subset \bigcup_{k=1}^m B(y_k, r_k) = U$$

and

$$\frac{\partial v_k}{\partial x_j}(x) = h_j(x), \quad \forall x \in B(y_k, r_k), \quad k = 1, \dots, m \text{ and } j = 1, \dots, n.$$

Moreover U is connected since $\partial\Omega$ is connected. Suppose that $A_{il} = B(y_i, r_i) \cap B(y_l, r_l) \neq \emptyset$. We have

$$\frac{\partial v_i}{\partial x_j}(x) = \frac{\partial v_l}{\partial x_j}(x), \quad \forall x \in A_{il}, \quad j = 1, \dots, n,$$

hence $v_i = v_l + c_{il}$ on A_{il} for some constant c_{il} . Define

$$v = \begin{cases} v_i & \text{on } B(y_i, r_i), \\ v_l + c_{il} & \text{on } B(y_l, r_l). \end{cases}$$

Then v is analytic on $B(y_i, r_i) \cup B(y_l, r_l)$ and

$$\frac{\partial v}{\partial x_j}(x) = h_j(x), \quad \forall x \in B(y_i, r_i) \cup B(y_l, r_l), \quad j = 1, \dots, n.$$

The lemma follows by repeating this procedure. \square

Lemma 8. *There exist an open connected neighborhood V of Ω and a function w analytic in V such that*

$$\frac{\partial w}{\partial x_j}(x) = h_j(x), \quad \forall x \in V, \quad j = 1, \dots, n.$$

Proof. With the notations of Lemma 7 let $y \in U \cap \Omega$ and let $r = d(y, \partial\Omega)$. Since h_j is analytic in Ω ,

$$h_j(x) = \sum_{\alpha \geq 0} \frac{1}{\alpha!} \frac{\partial^{|\alpha|} h_j}{\partial x^\alpha}(y) (x - y)^\alpha$$

for $x \in B(y, r)$ and $j = 1, \dots, n$. For $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ we define $w_\alpha \in \mathbb{R}$ as follows: $w_0 = v(y)$

$$w_{(\alpha_1, 0, \dots, 0)} = \frac{\partial^{\alpha_1-1} h_1}{\partial x_1^{\alpha_1-1}}(y) \quad \text{for } \alpha_1 \geq 1,$$

$$\vdots$$

$$w_{(0, \dots, 0, \alpha_n)} = \frac{\partial^{\alpha_n-1} h_n}{\partial x_n^{\alpha_n-1}}(y) \quad \text{for } \alpha_n \geq 1,$$

and, when there exist $\alpha_j, \alpha_k \geq 1$ for j and k such that $1 \leq j < k \leq n$

$$w_\alpha = \frac{\partial^{|\beta|+1} h_j}{\partial x_k \partial x^\beta}(y) = \frac{\partial^{|\beta|+1} h_k}{\partial x_j \partial x^\beta}(y),$$

where $\beta = (\alpha_1, \dots, \alpha_{j-1}, \alpha_j - 1, \alpha_{j+1}, \dots, \alpha_{k-1}, \alpha_k - 1, \alpha_{k+1}, \dots, \alpha_n)$ with the obvious modifications when $j = 1$ or $k = n$ or $k = j + 1$ and where the last equality is given by Lemma 7. Let again $\varepsilon_1 = (1, 0, \dots, 0), \dots, \varepsilon_n = (0, \dots, 0, 1)$. Since

$$\begin{aligned} h_j(x) &= \sum_{\alpha \geq 0} \frac{1}{\alpha!} w_{\alpha + \varepsilon_j} (x - y)^\alpha \\ &= \frac{\partial}{\partial x_j} \left(\sum_{\alpha \geq 0} \frac{1}{\alpha!} w_\alpha (x - y)^\alpha \right), \end{aligned}$$

for $x \in B(y, r)$ and $j = 1, \dots, n$, we define an analytic function w in $B(y, r)$ by

$$w(x) = \sum_{\alpha \geq 0} \frac{1}{\alpha!} w_\alpha (x - y)^\alpha, \quad x \in B(y, r),$$

such that

$$\frac{\partial w}{\partial x_j} = h_j \quad \text{on } U \cap B(y, r), \quad j = 1, \dots, n.$$

Since

$$\frac{\partial v}{\partial x_j} = h_j \quad \text{on } U \cap B(y, r), \quad j = 1, \dots, n,$$

we deduce that there exists a constant d such that $w = v + d$ on $U \cup B(y, r)$. Therefore we still have

$$\frac{\partial w}{\partial x_j} = h_j \quad \text{on } U \cup B(y, r), \quad j = 1, \dots, n,$$

and the lemma follows easily. \square

Lemma 9. *There exists a function u analytic in Ω such that*

$$\begin{aligned} \Delta u + 1 &= 0 \quad \text{in } \Omega, \\ u &= 0, \quad \frac{\partial u}{\partial \nu} = d \quad \text{on } \partial\Omega, \end{aligned}$$

where d denotes some constant.

Proof. The function w given by Lemma 8 satisfies

$$\Delta w = a \quad \text{in } \Omega$$

for some constant $a \in \mathbb{R}$ since

$$\frac{\partial \Delta w}{\partial x_j} = \Delta \frac{\partial w}{\partial x_j} = \Delta h_j = 0 \quad \text{in } \Omega, \quad j = 1, \dots, n.$$

For $1 \leq k \leq n-1$ we have

$$\begin{aligned} \frac{\partial}{\partial x_k}(w) &= \frac{\partial w}{\partial x_k} + \frac{\partial w}{\partial x_n} \frac{\partial \varphi}{\partial x_k} \\ &= \frac{1}{v_n}(h_k v_n - h_n v_k) = 0 \end{aligned}$$

on $\partial\Omega$ by (1). Since $\partial\Omega$ is connected, there exists a constant $b \in \mathbb{R}$ such that $w = b$ on $\partial\Omega$. If $a = 0$, then $w = b$ in Ω a contradiction to the fact that $h_j \neq 0$. Therefore $a \neq 0$. Now

$$u = -\frac{1}{a}w + \frac{b}{a}$$

satisfies Lemma 9. \square

Proof of Theorem 1 completed. Theorem 1 follows from Lemma 9 and Theorem 2. \square

3. Proof of the corollary

For every $j = 1, \dots, n$ let h_j be a harmonic function satisfying (1) with $c = C$. $C \neq 0$ by Remark 2. Let $h \in C^2(\Omega) \cap C^1(\overline{\Omega})$ be a harmonic function. From Green's identity we get

$$\int_{\partial\Omega} h_j \frac{\partial h}{\partial \nu} ds = \int_{\partial\Omega} h \frac{\partial h_j}{\partial \nu} ds \quad (23)$$

for $j = 1, \dots, n$. Then using (1), (3) and (23) we obtain

$$\int_{\partial\Omega} h \left(\frac{\partial h_j}{\partial \nu} + (1 + C(n-1)H) v_j \right) ds = 0 \quad (24)$$

for $j = 1, \dots, n$. Now choose h such that

$$h = \frac{\partial h_j}{\partial \nu} + (1 + C(n-1)H) v_j \quad \text{on } \partial\Omega.$$

Then (24) implies that (2) holds with $c = C$, and Theorem 1 completes the proof.

Acknowledgment

The author thanks the referee for her or his valuable comments.

References

- [1] A. Bennett, Symmetry in an overdetermined fourth order elliptic boundary value problem, *SIAM J. Math. Anal.* 17 (1986) 1354–1358.
- [2] H. Berestycki, L. Caffarelli, L. Nirenberg, Further qualitative properties for elliptic equations in unbounded domains, *Ann. Sc. Norm. Super. Pisa Cl. Sci.* (4) 25 (1–2) (1997) 69–94.
- [3] A. Farina, B. Kawohl, Remarks on an overdetermined boundary value problem, *Calc. Var. Partial Differential Equations* 31 (2008) 351–357.
- [4] A. Farina, E. Valdinoci, Flattening results for elliptic PDEs in unbounded domains with applications to overdetermined problems, *Arch. Ration. Mech. Anal.* 195 (3) (2010) 1025–1058.
- [5] I. Fragalà, F. Gazzola, Partially overdetermined elliptic boundary value problems, *J. Differential Equations* 245 (5) (2008) 1299–1322.
- [6] L. Hauswirth, F. Hélein, F. Pacard, On an overdetermined elliptic problem, *Pacific J. Math.* 250 (2) (2011) 319–334.
- [7] J. Prajapat, Serrin's result for domains with a corner or cusp, *Duke Math. J.* 91 (1998) 29–31.
- [8] J. Serrin, A symmetry problem in potential theory, *Arch. Ration. Mech. Anal.* 43 (1971) 304–318.
- [9] P. Sicbaldi, New extremal domains for the first eigenvalue of the Laplacian in flat tori, *Calc. Var. Partial Differential Equations* 37 (3–4) (2010) 329–344.
- [10] H.F. Weinberger, Remark on the preceding paper of Serrin, *Arch. Ration. Mech. Anal.* 43 (1971) 319–320.
- [11] N.B. Willms, G.M.L. Gladwell, D. Siegel, Symmetry theorems for some overdetermined boundary value problems on ring domains, *Z. Angew. Math. Phys.* 45 (4) (1994) 556–579.